


Subaru Ground-Layer AO Simulation




Shin Oya (Subaru Telescope)
Subaru Next Generation AO Working Group

2014/7/28 @ Mitaka

Basic idea of GLAO

- Corrects only turbulence close to the ground
- Improves seeing over wide-field of view

GLAO correction
(simulation)




Effective for wide field of view

Subaru GLAO configuration

$\phi = 15 \text{ arcmin}$
 $r = \phi/2$





LGS: 10 mag
TTFGS: 18mag

Subaru seeing:
-good: 0.56"
-moderate: 0.73"
-bad: 0.97"
(at 0.5 μm)



Seeing dependence of FWHM


@zenith



Stalibiy
≡ std / ave
GLAO: 1~7%
Seeing: 2~10%

GLAO: solid , Seeing: dotted; error bars shows standard deviation along time axis

Seeing dependence of Ensquared Energy





@zenith

Gain \equiv
GLAO / Seeing
 ~ 1.5 to 2

width: blue: 0.24", green: 0.36", red: 0.48"
GLAO: solid lines, Seeing: dotted line



Zenith angle dependency: GLAO / Seeing

FWHM



effective height
increases

seeing:
-good
-moderate
-bad




loss by 20% at 45° and 30% at 60°

Seasonal Variation of Seeing


Subaru IQ

<http://www.subaru.nao.ac.jp/Observing/Telescope/ImageQuality/Seeing/>



13N site, profile

<http://sitedata.tmt.org>





Characteristic months Sep (good) & Jan (bad).

Subaru AG

25%-ile	Sep (50%-ile)	50%-ile	Jan (50%-ile)	75%-ile
0.49"	0.54"	0.64"	0.72"	0.84"

Seasonal Variation of FWHM




@zenith

Variation
+/-30%
(w/o Jan)

GLAO: R: green, J: blue, H: magenta, K: red; Seeing: black
moderate: dotted line

FWHM ratio (GLAO/Seeing) Map




shift the map: late \leftarrow midnight \rightarrow early

Sky coverage

B-configuration

★:LGS

●: TTFGS



diameter of the dotted circle is 7.5'

preliminary

Number of TTFGS in each of 4 pink region

D. Simons, Gemini technical notes TN-PS-G0030, (1995).

$r(\text{in, out}) = r_{\text{circ}}$	TTFGS(R-band)	$b: 10 \sim 20 \text{ deg}$	$30 \sim 50 \text{ deg}$	$60 \sim 90 \text{ deg}$	
$(7', 8') = 1.6'$	< 18 mag	> 6.7	> 3.0	> 1.8	standard
	< 19 mag	> 10.7	> 4.8	> 2.8	1mag dim
$(7', 8.5') = 2.0'$	< 18mag	> 34.7	> 7.4	> 4.3	1' larger dia



Ground layer measurement at Subaru

Local ground-layer at Subaru?


On-going!!

- 70m below and leeward of the ridge (laminar flow?)
- fine resolution data for more detailed simulation

Luna Shabar (PTP) by Univ.BC
optical: 1 ~ 1000m




SNODAR by Univ. NSW
acoustic: 10 ~ 100m




RAVEN at Subaru



RAVEN is a Multi-Object AO (MOAO) demonstrator, yet has GLAO mode



Succeeded in the 1st light in May!!



RAVEN first light result



Summary

- Expected FWHM is 0.2" in the K-band under moderate seeing condition. Stability (std/ave) along time axis is same or better than seeing.
- Gain of ensquared energy is ~ 1.5 to 2
- Gain of FWHM decreased with zenith angle by 20% at 45deg and by 30% at 60deg
- Seasonal variation of FWHM is ~30% (except Jan)
- Even at galactic pole, expected number of an 18mag star in each of 4 TTFGS of $\phi 15'$ case is > 1.
- Ground-layer evaluation at Subaru is on-going.
- FWHM < 0.2" in the H-band was actually achieved by RAVEN GLAO mode.